



A POWH Ophthalmology / CFEH Collaboration

GP REFERRAL: GLAUCOMA MANAGEMENT CLINIC

General Practitioner to FAX to (02) 8115 0799

Patient Contact Details	Assistance Required
Title: Dr Mr Mrs Miss Ms	Other: Mobility: Wheelchair Other:
First Name:	Language Interpreter: ☐ Yes ☐ No
	If yes, specify language:
Surname:	Hearing Interpreter:
Mailing Address:	
Suburb:	NOTE: Urgent referrals should not be sent to the Centre
Postcode: State:_	
Reason for Referral	
☐ Is under treatment for (non-advanced*) glaucoma but not currently under regular optometrist / ophthalmologist care	
☐ Is under treatment for (non-advanced*) glaucoma and due to financial difficulty has requested an alternative option	
☐ Is a glaucoma suspect [#] needing further evaluation and not currently under regular review	
□ Other (glaucoma)	
Medical and Ocular History (please attach additional information including medications and previous treatment as appropriate)	
Patient's previous or current optometrist / ophthalmologist (if applicable)	
Ophthalmologist's Name:	Practice location:
Optometrist's Name	Practice location:
Referring Practitioner Details	
Name:	Practice Name and Address:
Medicare Provider No:	
Date://	Signature:





A POWH Ophthalmology / CFEH Collaboration

Please note that:

- Where patients have a family history of glaucoma or other risk factors and you wish to have them assessed, we recommend they are initially referred to a local optometrist or ophthalmologist.
- All tests conducted at CFEH are routine clinical tests and do not involve experimental procedures.
- UNSW optometry students may observe the consultation and assist with visual field testing and other imaging as needed
- Patients should **not** be referred for acute and /or urgent conditions including angle closure glaucoma. On the rare occasion that the patient presents with a condition that requires urgent treatment, appropriate clinical management will be formulated in consultation with the referring practitioner where possible
- The GMC is a satellite clinic of the SESLHD with the consultants employed by the SESLHD

Glaucoma suspects include patients with no visual field defects but who have significant risk factors for developing the condition including: a strong family history (first generation), high myopia, history of eye trauma, regular corticosteroid usage, thin corneas and pseudo-exfoliation or pigment dispersion syndrome.

^{*} Patients with advanced glaucoma are defined as having definite optic nerve head pathology and repeatable visual field loss over 12 dB and/or within 10 degrees of fixation (ie marked and/or central loss of visual field)